

UT-800
December 1997

BPS Condition of String Junction from M Theory

YUTAKA MATSUO*, KAZUMI OKUYAMA[†]

Department of Physics, Faculty of Science
Tokyo University
Bunkyo-ku, Hongo 7-3-1, Tokyo 113, Japan

Abstract

We give a simple derivation of BPS condition of string junction from M theory

hep-th/9712070

* e-mail address : matsuo@phys.s.u-tokyo.ac.jp

[†] e-mail address : okuyama@hep-th.phys.s.u-tokyo.ac.jp

Recently BPS condition of the (p, q) -string junction¹ conjectured in [2] was derived in the linearized approximation by Dasgupta and Mukhi [3]. Such a configuration seems physically relevant in the description of the exceptional groups from open string theory [4]. It might be also important in the understanding of manifestly $SL(2, Z)$ invariant formulation in type IIB superstring theory [5]. Subsequently further study was made on the string network [6] and BPS dynamics [7].

BPS condition in [3] may be summarized as follows. We are considering a junction of three (p_i, q_i) strings ($i = 1, 2, 3$) (Fig.1) stretching in $\vec{n}_i$ directions in X^1 - X^2 plane ($|n_i| = 1$). To keep supersymmetry, we need to impose




Figure 1: String Junction

1. the string charges should be conserved [2],

$$\sum_{i=1}^3 p_i = \sum_{i=1}^3 q_i = 0. \quad (1)$$

2. the forces from each string tensions should be balanced [3],

$$\sum_{i=1}^3 T_i \vec{n}_i = 0, \quad T_i = |p_i \tau + q_i|, \quad \tau = \chi + i e^{-\phi}. \quad (2)$$

¹Similar (p, q) -fivebrane junction is considered in ref.[1]

In this short note, we will attempt to give a simplified derivation of BPS condition from M-theory viewpoint[8]. BPS configuration of M-theory (or more exactly supermembrane) was discussed by using κ -symmetry [9] when target space is Calabi-Yau manifold. For the toroidal target space, it was examined in [10] in the light cone gauge and found that the supersymmetric configurations are essentially restricted to double dimensional reduction of the string BPS configuration, namely holomorphic map in two directions and trivial winding in third direction.² We show that the requirement of holomorphy directly gives the BPS condition discussed above.

Recall that the (p, q) string is described in M theory as a membrane winding q (resp. p) times in X^{11} (resp. X^9) direction. Namely if we parametrize the spatial part of the world volume of membrane by σ_1, σ_2 where the winding direction described by σ_2 , the boundary condition is given as,

$$W(\sigma_1, \sigma_2 + 2\pi) = W(\sigma_1, \sigma_2) + p\tau + q, \quad W = X^{11} + iX^9. \quad (3)$$

The string junction is then described as a “pants” diagram (Fig.2 left). In the infinite leg limit, this diagram is holomorphically equivalent to a sphere with three punctures (Fig.2 right) We take a holomorphic coordinate ζ on




Figure 2: Pants diagram

this sphere where the three legs are attached to points ζ_i ($i = 1, 2, 3$). From the mapping in Fig.2, the asymptotic behaviour of the embedding into target space near each singularity should look like,

$$\begin{aligned} Z &\equiv X^1 + iX^2 \sim \mathcal{S}_i \tilde{\sigma}_1^{(i)}, \\ W &\sim \mathcal{T}_i \tilde{\sigma}_2^{(i)}, \end{aligned} \quad (4)$$

² We note that similar construction of BPS state was used by Witten[11] in the derivation of Seiberg-Witten curve from fivebrane configuration.

where $\mathcal{T}_i = p_i\tau + q_i$, $\arg(\mathcal{S}_i) = \theta_i$ with

$$\vec{n}_i = \begin{pmatrix} \cos \theta_i \\ \sin \theta_i \end{pmatrix}, \quad \log(\zeta - \zeta_i) = \tilde{\sigma}_1^{(i)} + i\tilde{\sigma}_2^{(i)}.$$

This embedding does not look like holomorphic as long as we take W and Z as holomorphic variables. Our strategy is to replace it by an appropriate complex structure in this four dimensional target space such that the above embedding becomes holomorphic. This condition actually gives over-constrained system of equations and the BPS conditions (1), (2) becomes conditions where they have solutions.

We write two complex numbers W, Z as four dimensional vector, $\vec{Y} = (X^{11}, X^9, X^1, X^2)^t$ and let J be a complex structure described by four by four matrix satisfying,

$$J^2 = -I_4, \quad J^t \cdot J = I_4. \quad (5)$$

Cauchy-Riemann relation can be written as,

$$\partial_{\sigma_1} \vec{Y} = \partial_{\sigma_2} (J\vec{Y}), \quad (6)$$

where we choose a particular holomorphic coordinate of the original Riemann sphere $\zeta = \sigma_1 + i\sigma_2$. At the asymptotic region, we may replace them by $\tilde{\sigma}_{\{1,2\}}^{(i)}$. (6) then gives,

$$\begin{pmatrix} 0 \\ 0 \\ S_1^{(i)} \\ S_2^{(i)} \end{pmatrix} = J \cdot \begin{pmatrix} T_1^{(i)} \\ T_2^{(i)} \\ 0 \\ 0 \end{pmatrix}, \quad \mathcal{S}^{(i)} = S_1^{(i)} + iS_2^{(i)}, \quad \mathcal{T}^{(i)} = T_1^{(i)} + iT_2^{(i)}. \quad (7)$$

One may write J as

$$J = \begin{pmatrix} 0 & m_1 \\ m_2 & 0 \end{pmatrix}. \quad (8)$$

The conditions (5) reduce to following conditions on m ,

$$m_1 \cdot m_2 = -I_2, \quad m_i \cdot m_i^t = I_2.$$

Since matrix m now belongs to $O(2)$, one can immediately recognize that Cauchy-Riemann relation gives

$$S_1^{(i)} + iS_2^{(i)} = e^{i\varphi} (T_1^{(i)} \pm iT_2^{(i)}). \quad (9)$$

Namely, up to rotation and parity in X^1 - X^2 plane, the spacial orientation of each string is determined from its charge (p, q) . This condition seems stronger than the requirement (2). However, as observed in [6] they are equivalent. Finally, the charge conservation (1) comes from the integration,

$$\sum_i \int_{C_i} d(\vec{Y} + iJ\vec{Y}) = 0. \quad (10)$$

where C_i is a circle which surrounds the point $\zeta = \zeta_i$.

Although we used only the asymptotic behaviour of the embedding, it is obvious that such embedding exists globally from the very nature of holomorphy. This completes our derivation of BPS condition. At this point, it is obvious to extend our analysis to n -string junction or “string network” discussed by Sen [6] (the holomorphic map of a Riemann surface with n punctures).

After we finished our computation, we saw a paper [12] which discuss the string junction in the similar line.

References

- [1] O. Aharony, A. Hanany and B. Kol, “Webs of (p,q) 5-branes, Five Dimensional Field Theories and Grid Diagrams ”, hep-th/9710116.
- [2] J. H. Schwarz, Nucl. Phys. Proc. Suppl. 55B (1997) 1, hep-th/9607201.
- [3] K. Dasgupta and S. Mukhi, “BPS Nature of 3-String Junctions”, hep-th/9711094.
- [4] M. R. Gaberdiel and B. Zwiebach, “Exceptional Groups from Open Strings”, hep-th/9709013.
 See also,
 K. Dasgupta and S. Mukhi, Phys. Lett. B385 (1996) 125, hep-th/9606044.
 A. Johansen, Phys. Lett. B395 (1997) 36, hep-th/9608186.
- [5] See for example, P. K. Townsend, “Membrane tension and manifest IIB S-duality”, hep-th/9705160.
- [6] A. Sen, “String Network”, hep-th/9711130.

- [7] S.-J. Rey and J.-T. Yee, “BPS Dynamics of Triple (p,q) String Junction”, hep-th/9711202.
- [8] O. Aharony, J. Sonnenschein and S. Yankielowicz, Nucl. Phys. B474 (1996) 309, hep-th/9603009.
- [9] K. Becker, M. Becker and A. Strominger, “Fivebranes, Membranes and Non-Perturbative String Theory”, hep-th/9507158.
- [10] K. Ezawa, Y. Matsuo, K. Murakami, hep-th/9706002, hep-th/9707200.
- [11] E. Witten, “Solutions of Four-Dimensional Field Theories Via M Theory”, hep-th/9703166.
- [12] M. Krogh and S. Lee, “String Network from M-theory”, hep-th/9712050.